

# Human recombinant RAGE (Receptor for advanced glycation endproducts) protein, AF

Catalog Number: PROTQ15109-2

## Overview

Product Name	Human recombinant RAGE (Receptor for advanced glycation endproducts) protein, AF
Description	Receptor for advanced glycation endproducts (RAGE) is a 35 kDa transmembrane receptor of the immunoglobulin super family. The mature RAGE has three main parts, consisting of extracellular, transmembrane, and cytosolic regions. A central mechanism by which ligand-RAGE interaction mediates cell stress and upregulates inflammatory pathways is via activation of signal transduction pathways.
Size	5ug,20ug,100ug
Tag	His Tag (C-term)
Form	Lyophilized
Source	Escherichia coli
Formulation	The protein was lyophilized from a 0.2 um filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.

## Storage

Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

## Purity

>98% as determined by SDS-PAGE.

## Molecular weight

The protein has a calculated MW of 30.94 kDa. The protein migrates as 40 kDa under reducing condition (SDS-PAGE analysis).

## Activity

Testing in process

## Endotoxin

<0.1 EU per 1 ug of the protein by the LAL method.

### Amino Acid Sequence

MAQNITARIGEPLVLKCKGAPKKPPQRLEWKLNTGRTEAWKVLSPQGGGPWDSVARVLPNGSLFLPAVGIQDEGIFRCQAMNRNGKETKSNYRVRVY  
QIPGKPEIVDSASELTAGVPNKVGTVCVSEGSYPAGTLSWHLDGKPLVPNEKGVSVKEQTRRHPETGLFTLQSELMVTPARGGDPPTFSCSFSPGLPRHR  
ALRTAPIQPRVWEPVPLEEVQLVVEPEGGAVAPGGTVTLTCEVPAQPSPQIHWMKDGVPLPLPPSPVLILPEIGPQDQGTYS with polyhistidine tag at  
the C-terminus.

### Reconstitution

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H<sub>2</sub>O to a concentration not less than 100 ug/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

## Usage

---

Boster's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

## Submit a product review to Biocompare.com

---

Submit a review of this product to Biocompare.com to receive a \$20 Amazon giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Human recombinant RAGE (Receptor for advanced glycation endproducts) protein, AF

For Research Use Only. Not for use in diagnostic procedures.