

## Human recombinant IL-30 (Interleukin-30) protein, AF

Catalog Number: PROTQ8NEV9-1

### Overview

Product Name	Human recombinant IL-30 (Interleukin-30) protein, AF
Description	Interleukin 30 (IL-30) is a 27 kDa cytokine with 216 amino acid residues. IL-30, also known as IL-27A, forms a complex named IL-27 with Ebi3 and is secreted from cells of myeloid origin, such as monocytes, macrophages, and dendritic cells. IL-30/Ebi3 complex mediates T-cell immunity, promoting Th1 differentiation, inhibiting Th2 differentiation, and inhibiting Th17 differentiation. IL-30 also can be secreted independently of Ebi3 and modulate bacteria-induced oxidative burst and cytokine production.
Size	5ug,20ug,100ug
Tag	His Tag (C-term)
Form	Lyophilized
Source	Escherichia coli
Formulation	The protein was lyophilized from a 0.2 um filtered solution containing 1X PBS, pH 8.0. If you have any concerns or special requirements, please confirm with us.

### Storage

Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

### Purity

>98% as determined by SDS-PAGE.

### Molecular weight

The protein has a calculated MW of 25.47 kDa. The protein migrates as 28 kDa under reducing condition (SDS-PAGE analysis).

### Activity

Testing in process

### Endotoxin

<0.1 EU per 1 ug of the protein by the LAL method.

### Amino Acid Sequence

MFPRPPGRPQLSLQELRREFTVSLHLARKLLSEVRGQAHRF AESHLPGVNLYLLPLGEQLPDVSLTFQAWRR LSDPERLCFISTTLQPFHALLGGLGTQGR  
WTNMERMQLWAMRLDLRDLQRHLRFQVLAAGFNLPEEEEEEEEEEEERKGLLPGALGSALQGPAQVSWPQLLSTYRLLHSLELVLSRAVRELLLLSKA  
GHSVWPLGFPTLSPQP with polyhistidine tag at the C-terminus.

### Reconstitution

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H<sub>2</sub>O to a concentration not less than 100 ug/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

## Usage

---

Boster's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

## Submit a product review to Biocompare.com

---

Submit a review of this product to Biocompare.com to receive a \$20 Amazon giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Human recombinant IL-30 (Interleukin-30) protein, AF

For Research Use Only. Not for use in diagnostic procedures.