BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Mouse recombinant LIGHT protein, AF

Catalog Number: PROTQ9QYH9-2

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a $20 Amazon giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Mouse recombinant LIGHT protein, AF
Description	LIGHT, also known as TNFSF14 is a member of the TNF superfamily that produced by multiple immune cells such as activated T cells and immature dendritic cells (DCs). Mouse LIGHT shares 79% sequence homology with human LIGHT. LIGHT is a 29 kDa type II transmembrane protein, which serves as a ligand for both lymphotoxin beta receptor (LTbetaR) and TNFSF signaling receptors (TNFRSF14/HVEM). Besides, LIGHT can amplify NF-kB signaling pathway in T cells when presencing anti-CD3 antibody treatment. Additionally, LIGHT is able to stimulate T cell proliferation and IFN expression via interacting with TNFRS13 /HVEM.
Size	5ug,20ug,100ug
Tag	His Tag (C-term)
Form	Lyophilized
Source	Escherichia coli
Formulation	The protein was lyophilized from a 0.2 um filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.

Storage

Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

Purity

>98% as determined by SDS-PAGE.

Molecular weight

The protein has a calculated MW of 20.92 kDa. The protein migrates about 17 kDa under reducing condition (SDS-PAGE analysis).

Activity

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Measure by its ability to induce cytotoxicity in HT-29 cells. The ED_{50} for this effect is <2 ug/mL.

Endotoxin

<0.1 EU per 1 ug of the protein by the LAL method.

Amino Acid Sequence

MRLHQRLGDIVAHLPDGGKGSWEKLIQDQRSHQANPAAHLTGANASLIGIGGPLLWETRLGLAFLRGLTYHDGALVTMEPGYYYVYSKVQLSGVGCPQ GLANGLPITHGLYKRTSRYPKELELLVSRRSPCGRANSSRVWWDSSFLGGVVHLEAGEEVVVRVPGNRLVRPRDGTRSYFGAFMV with polyhistidine tag at the C-terminus.

Reconstitution

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H_2O to a concentration not less than 100 ug/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

BOSTER
antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Usage

Boster's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Mouse recombinant LIGHT protein, AF

For Research Use Only. Not for use in diagnostic procedures.