

## Human recombinant AITRL (Activation-induced TNFR member ligand) protein, AF

Catalog Number: PROTQ9UNG2-7

### Overview

Product Name	Human recombinant AITRL (Activation-induced TNFR member ligand) protein, AF
Description	AITRL also known as TNFSF18, GITRL and TL6, belonging to TNF superfamily. AITRL is a 20.3 kDa protein containing 177 residues that implicates in regulating immune systems. AITRL contributes to tumor suppression and the development of autoimmune diseases via interacting with TNFRSF18/AITR/GITR. Besides, AITRL-GITR signaling has been demonstrated to enhance phosphorylation of STAT1 and up-regulate expression of VCAM1 and ICAM1 in endothelial cells. Furthermore, AITRL can also facilitate the adhesion and transmigration of leukocytes to endothelial cells.
Size	5ug,20ug,100ug
Tag	His Tag (C-term)
Form	Lyophilized
Source	Escherichia coli
Formulation	The protein was lyophilized from a 0.2 um filtered solution containing 1X PBS, pH 7.4. If you have any concerns or special requirements, please confirm with us.

### Storage

Lyophilized protein should be stored at -20°C for 1 year. Upon reconstitution, store at 2°C to 8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10%FBS, 5%HSA or 5% trehalose solution), protein aliquots should be stored at -20°C or -80°C for 3-6 months. Avoid repeated freeze/thaw cycles.

### Purity

>98% as determined by SDS-PAGE.

### Molecular weight

The protein has a calculated MW of 15.34 kDa. The protein migrates as 11 kDa under reducing condition (SDS-PAGE analysis).

### Activity

Measure by its ability to induce IL-8 secretion in human PBMC using a concentration range of 5-200 ng /mL. Note: Result may vary from different PBMC donors.

### Endotoxin

<0.1 EU per 1 ug of the protein by the LAL method.

### **Amino Acid Sequence**

MQLETAKEPCMAKFGPLPSKWQMASSEPPCVNKVSDWKLEILQNGLYLIYGQVAPNANYNDVAPFEVRLYKNKDMIQTLTNKSKIQNVGGTYELHVG  
DTIDLIFNSEHQVLKNNTYWGII LIANPQEI with polyhistidine tag at the C-terminus.

### **Reconstitution**

Centrifuge at 3000 rpm for 5 mins before opening. It is recommended to reconstitute the lyophilized protein in sterile H<sub>2</sub>O to a concentration not less than 100 ug/mL and incubate the stock solution at room temperature for at least 20 mins to ensure sufficient re-dissolved. Do Not Vortex!  
Vigorous shaking may impair the biological activity of the protein.

## Usage

---

Boster's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

## Submit a product review to Biocompare.com

---

Submit a review of this product to Biocompare.com to receive a \$20 Amazon giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Human recombinant AITRL (Activation-induced TNFR member ligand) protein, AF

For Research Use Only. Not for use in diagnostic procedures.