



## **ISRE Luciferase Reporter- RAW264.7 Cell Line**

**Catalog number: RC1041**

This package insert must be read in its entirety before using this product.  
For research use only. Not for use in diagnostic procedures.

### **ISRE Luciferase Reporter-RAW264.7 Cell Line**

**Catalog Number:** RC1041, **Storage:** Immediately upon receipt, store in liquid nitrogen. (Ship on dry ice.)

**Contents:** Each vial contains  $2 \sim 3 \times 10^6$  cells in 1 ml of 90% FBS + 10% DMSO.

**Description:** The ISRE Luciferase Reporter cell line is a stably transfected RAW264.7 cell line which expresses Renilla luciferase [reporter gene](#) under the transcriptional control of the Interferon-Stimulated Response Element (ISRE), so that the cell line is designed to monitor the JAK/STAT signaling pathway activity. This cell line can be activated by type I IFNs as well as certain Toll like receptor ligands capable of induction of IRFs such as TLR3 ligand-poly (I:C). Functional activity of the cell line has been validated by poly (I:C) (Figure 1).

**Applications:** Functional Assay

**Application Notes:** Functional Assay, detecting the transcriptional activity of ISRE

**Application Details:**

## Application:

Monitor the JAK/STAT signaling pathway activity. Screen for activators or inhibitors of the JAK/STAT signaling pathway.

## Culture conditions:

Cells should be grown at 37°C with 5% CO<sub>2</sub> using DMEM medium supplemented with 10% FBS and 1% Pen/Strep, plus 3 µg/ml of Puromycin. It is recommended to quickly thaw the frozen cells upon receipt or from liquid nitrogen in a 37°C water-bath, transfer to a tube containing 10 ml of growth medium without Puromycin, spin down cells, resuspend cells in pre-warmed growth medium without Puromycin, transfer resuspended cells to T25 flask and culture in 37°C-CO<sub>2</sub> incubator. Leave the T25 flask in the incubator for 1~2 days without disturbing or changing the medium until cells completely recover viability and become adherent. Once cells are over 90% adherent, remove growth medium and passage the cells through trypsinization and centrifugation. At first passage, switch to growth medium containing Puromycin. Cells should be split before they reach complete confluence. To passage the cells, detach cells from culture vessel with Trypsin/EDTA, add complete growth medium and transfer to a tube, spin down cells, resuspend cells and seed appropriate aliquots of cells suspension into new culture vessels. Subcultivation ration = 1:10 to 1:20 weekly.

## Functional validation:

A. Response of ISRE RAW264.7 cells to poly (I:C). 1. Harvest ISRE RAW264.7 cells and seed cells into a white solid-bottom 96-well microplate in 100 µl of growth medium at  $8.5 \times 10^4$  cells/well. 2. Incubate cells at 37°C in a CO<sub>2</sub> incubator for overnight. 3. The next day, stimulate cells with various concentrations of poly (I:C). 4. Incubate at 37°C in a CO<sub>2</sub> incubator for 6-16 hours. 5. Add 50 µl of luciferase assay reagent per well. 6. Incubate at room temperature for 1-5 minutes and measure luminescence using a microplate luminometer.

## ISRE Luciferase Reporter-RAW264.7 Cell Line (RC1041) Images




Fig-1: Induction of ISRE activity by poly(I:C) in ISRE RAW264.7 cells.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



ISRE Luciferase Reporter-RAW264.7 Cell Line

For Research Use Only. Not for use in diagnostic procedures.