

Anti-Interleukin-4 IL4 Antibody Picoband®

Catalog Number: RP1009

About IL4

Interleukin-4 (IL-4), also known as a B-cell stimulatory factor 1 (BSF1), is an immunomodulatory cytokine, which can inhibit the growth of tumour cells.¹ The human cDNA contains a single open reading frame encoding a protein of 153 amino acids, including a putative signal peptide. IL-4 may act as an autocrine growth factor in pancreatic cancer cells and also give rise to the possibility that cancer-derived IL-4 may suppress cancer-directed immunosurveillance in vivo in addition to its growth-promoting effects, thereby facilitating pancreatic tumor growth and metastasis.¹ The mouse and human genes and their protein products show structural and functional similarities. The human IL-4 gene, which occurs as a single copy in the haploid genome, is mapped on chromosome 5.² The standard product used in this kit is recombinant human IL-4, consisting of 130 amino acids with the molecular mass of 14KDa.

Overview

Product Name	Anti-Interleukin-4 IL4 Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-Interleukin-4 IL4 Antibody catalog # RP1009. Tested in ELISA, IHC, WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Na ₃ . Carrier free (No BSA) form available in stock. If you want this antibody carrier free please specify "Carrier Free" or "No BSA" in your order note.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P05112

Technical Details


Immunogen	E. coli-derived human IL-4 recombinant protein (Position: H25-S153).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins

Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human ELISA, 0.1-0.5ug/ml, - Western blot, 0.1-0.5ug/ml, Human


Anti-Interleukin-4 IL4 Antibody Picoband® (RP1009) Images



Figure. Western blot analysis of IL-4 using anti-IL-4 antibody (RP1009). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane: Recombinant Human IL-4 Protein 0.5ng, After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-IL-4 antigen affinity purified polyclonal antibody (Catalog # RP1009) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for IL-4 at approximately 14KD. The expected band size for IL-4 is at 14KD.




Protective effect of M2 macrophages induced by hUCMSCs-EVs on OA chondrocytes in vitro. A IL-1beta-induced OA chondrocytes were co-cultured with the supernatant of M2 macrophages (M2S) induced by hUCMSCs-EVs, or platelet-rich plasma (PRP) for 48 h, relative mRNA expression of the key genes TNF-alpha, MMP13, SOX9, and ACAN was measured by quantitative RT-PCR analysis; the experiment was performed triplicate; *p<0.05, **p<0.01, ***p<0.001. B Western blot was performed to evaluate the expression of TNF-alpha, MMP13, and IL-4 proteins in PBS, M2S, or PRP-treated OA chondrocytes; GAPDH was employed as the loading control; *p<0.05, **p<0.01, ***p<0.001. C The influence of M2S or PRP on the viability of chondrocytes was detected by the cell live/death experiment; green represents live cells while red represents dead cells; Scale bar: 1 mm Index in PubMed under a CC BY license. PMID: 35057811




Serum levels of IP-10, IFN-gamma, IL-4, and TGF-beta1 as well as the IFN-gamma/IL-4 ratio in chronic hepatitis B patients with or without fibrosis. The levels of serum IP-10, IFN-gamma, IL-4, and TGF-beta1 in CHB patients with or without liver fibrosis were determined by ELISA, and the IFN-gamma/IL-4 ratio was calculated. ♦♦♦ Differs from controls (the F0 group), P<0.05; ★★ ★ differs from mild or minimal fibrosis (the F1-2 group), P<0.05; ▲▲ ▲ differs from moderate fibrosis (the F3-4 group), P<0.05. (a) IP-10; (b) IFN-gamma; (c) IL-4; (d) TGF-beta1; (e) the IFN-gamma/IL-4 ratio. Index in PubMed under a CC BY license. PMID: 28067328


Statistical analysis of the correlation between the serum IP-10 level or the IFN-gamma/IL-4 ratio with liver fibrosis




among chronic hepatitis B patients. Spearman's correlation analysis of the association between (a) IP-10; (b) IFN-gamma; (c) the IFN-gamma/IL-4 ratio and TGF-beta1. Spearman's correlation analysis of the association between serum (d) IFN-gamma; (e) IL-4; (f) the IFN-gamma/IL-4 ratio and IP-10. Index in PubMed under a CC BY license. PMID: 28067328



ROC curve analysis for evaluating the sensitivity and specificity of the IP-10 level, IFN-gamma/IL-4 ratio, or their combination to predict significant fibrosis among CHB patients. (a) ROC curve analysis for serum IP-10 (with the cut-off value of 300 pg/mL), the serum IFN-gamma/IL-4 ratio (with the cut off value of 1.8), and the combination of IP-10 and the IFN-gamma/IL-4 ratio; (b) Specificity and sensitivity for IP-10, the IFN-gamma/IL-4 ratio, and their combination to predict significant liver fibrosis among patients with CHB. Index in PubMed under a CC BY license. PMID: 28067328



Intrahepatic mRNA levels of IP-10, IFN-gamma, and IL-4 in chronic hepatitis B patients with or without fibrosis. Real-time qRT-PCR was conducted to quantify the mRNA levels of intrahepatic IP-10, IFN-gamma, and IL-4 in the CHB patients without or with fibrosis as described in the Materials and Methods section. The relative mRNA levels of intrahepatic IP-10, IFN-gamma, and IL-4 were calculated by comparative Ct analysis after normalization for the quantity of GAPDH in the same samples and were represented as $2^{-\Delta\Delta Ct}$ values for controls (the F0 group), which were set equal 1. ♦♦♦ Differs from controls (the F0 group), $P < 0.05$; ★★ ★ differs from mild or minimal fibrosis (the F1-2 group), $P < 0.05$; ▲▲ ▲ differs from moderate fibrosis (the F3-4 group), $P < 0.05$. (a) IP-10; (b) IFN-gamma; (c) IL-4. Index in PubMed under a CC BY license. PMID: 28067328



Intrahepatic protein expression of IP-10, IFN-gamma, IL-4, TGF-beta1, and alpha-SMA as well as the IFN-gamma/IL-4 ratio in chronic hepatitis B patients with or without fibrosis. The protein expression of intrahepatic (a, b, c, and d) IP-10, (e, f, g, and h) IFN-gamma, and (i, g, k, and l) IL-4. In addition, the protein levels of intrahepatic IP-10, IFN-gamma, IL-4, TGF-beta1, and alpha-SMA were quantified based on the value of integrated optical density (IOD) and represented as histograms, from which the IFN-gamma/IL-4 ratio was calculated. ♦♦♦ Differs from controls (the F0 group), $P < 0.05$; ★★ ★ differs from mild or minimal fibrosis (the F1-2 group), $P < 0.05$; ▲▲ ▲ differs from moderate fibrosis (the F3-4 group), $P < 0.05$. Index in PubMed under a CC BY license. PMID: 28067328

4 Publications Citing This Product

1. PubMed ID: 23065188, Wang Yz, Feng Xg, Shi Qg, Hao Yl, Yang Y, Zhang Am, Kong Qx. Inflammation. 2013 Apr;36(2):337-45. Doi: 10.1007/S10753-012-9551-5. Silencing Of Mir155 Promotes The Production Of Inflammatory Mediators In Guillain-Barr?? Syndrome In Vitro.

2. PubMed ID: 22387551, Upregulated protein arginine methyltransferase 1 by IL-4 increases eotaxin-1 expression in airway epithelial cells and participates in antigen-induced pulmonary %u2026

3. PubMed ID: 25483698, Yan X, Wang D, Liang F, Fu L, Guo C. Hum Vaccin Immunother. 2014;10(12):3491-8. Doi: 10.4161/Hv.36084. Hpv16L1-Attenuated Shigella Recombinant Vaccine Induced Strong Vaginal And Systemic Immune Responses In Guinea Pig Model.

Visit bosterbio.com/anti-human-il-4-antibody-rp1009-boster.html to see all 4 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Interleukin-4 IL4 Antibody

For Research Use Only. Not for use in diagnostic procedures.