

Anti-Interleukin-7 IL7 Antibody Picoband®

Catalog Number: RP1013

About IL7

IL-7, Interleukin 7, is a protein that in humans is encoded by the IL-7 gene. IL-7 a hematopoietic growth factor secreted by stromal cells in the red marrow and thymus. It is also produced by keratinocytes, dendritic cells, hepatocytes, neurons, and epithelial cells but is not produced by lymphocytes. By the combination of approaches, IL-7 is located on 8q12-q13. IL-7 is critical for early T-cell development and homeostasis of naive and memory CD8-positive T cells. It signals through the IL-7 receptor complex, which consists of the IL7R-alpha chain and the common IL2R-gamma chain is critical for early T-cell development and homeostasis of naive and memory CD8 -positive T cells. It signals through the IL-7 receptor complex, which consists of the IL7R-alpha chain and the common IL2R-gamma chain.

Overview

Product Name	Anti-Interleukin-7 IL7 Antibody Picoband®
Reactive Species	Human
Description	Boster Bio Anti-Interleukin-7 IL7 Antibody catalog # RP1013. Tested in ELISA, WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ . Carrier free (No BSA) form available in stock. If you want this antibody carrier free please specify "Carrier Free" or "No BSA" in your order note.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P13232

Technical Details


Immunogen	E. coli-derived human IL-7 recombinant protein (Position: D26-H177).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG

Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Human, - ELISA, 0.1-0.5ug/ml, -

Anti-Interleukin-7 IL7 Antibody Picoband® (RP1013) Images



Anti-human IL7 antibody, RP1013, Western blotting
Lane 1: Recombinant Human IL-7 Protein 10ng
Lane 2: Recombinant Human IL-7 Protein 5ng
Lane 3: Recombinant Human IL-7 Protein 2



Western blot analysis of IL-7 using anti-IL-7 antibody (RP1013). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human PC-3 whole cell lysates, Lane 2: human Hela whole cell lysates, After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-IL-7 antigen affinity purified polyclonal antibody (Catalog # RP1013) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for IL-7 at approximately 20KD. The expected band size for IL-7 is at 20KD.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Interleukin-7 IL7 Antibody

For Research Use Only. Not for use in diagnostic procedures.