

## Anti-Interleukin-10 IL10 Antibody Picoband®

Catalog Number: RP1016

### About IL10

Interleukin-10 (IL-10 or IL10), also known as human cytokine synthesis inhibitory factor (CSIF), is an anti-inflammatory cytokine. In humans IL-10 is encoded by the IL10 gene. It is capable of inhibiting synthesis of pro-inflammatory cytokines like IFN-gamma, IL-2, IL-3, TNFalpha and GM-CSF made by cells such as macrophages and regulatory T-cells. IL-10 also displays potent abilities to suppress the antigen presentation capacity of antigen presenting cells. Kim et al. (1992) showed that the mouse IL 10 gene contains 5 exons and spans about 5.2 kb of genomic DNA. Eskdale et al. (1997) mapped the IL10 gene to the junction between 1q31 and 1q32.

### Overview


Product Name	Anti-Interleukin-10 IL10 Antibody Picoband®
Reactive Species	Rat
Description	Boster Bio Anti-Interleukin-10 IL10 Antibody catalog # RP1016. Tested in ELISA, IHC, WB applications. This antibody reacts with Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	ELISA, IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P29456

### Technical Details


Immunogen	E. coli-derived rat IL-10 recombinant protein (Position: S19-N178).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml, Rat Immunohistochemistry (Paraffin-embedded Section), 2-5ug/ml, Rat ELISA, 0.1-0.5ug/ml, -

## Anti-Interleukin-10 IL10 Antibody Picoband® (RP1016) Images



Western blot analysis of IL10 using anti-IL10 antibody (RP1016). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. Lane 1: recombinant rat IL10 protein 10 ng, Lane 2: recombinant rat IL10 protein 5 ng, Lane 3: recombinant rat IL10 protein 1 ng. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-IL10 antigen affinity purified polyclonal antibody (Catalog # RP1016) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for IL10 at approximately 17 kDa.



IHC analysis of IL10 using anti-IL10 antibody (RP1016). IL10 was detected in a paraffin-embedded section of rat spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-IL10 Antibody (RP1016) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

## 20 Publications Citing This Product

1. PubMed ID: PMID:27347328, Pentoxifylline inhibits pulmonary inflammation induced by infrarenal aortic cross-clamping dependent of adenosine receptor A2A
2. PubMed ID: 10.1177/1744806918788681, Nuclear factor kappa B regulated monocyte chemoattractant protein-1/chemokine CC motif receptor-2 expressing in spinal cord contributes to the maintenance of cancer-induced bone pain in rats:
3. PubMed ID: 10.1002/cbf.1523, Inhibitory effect of mesenchymal stem cells on lymphocyte proliferation

Visit [bosterbio.com/anti-rat-il-10-antibody-rp1016-boster.html](http://bosterbio.com/anti-rat-il-10-antibody-rp1016-boster.html) to see all 20 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Interleukin-10 IL10 Antibody

For Research Use Only. Not for use in diagnostic procedures.