

Anti-Interleukin-18 IL18 Antibody Picoband® APC Conjugated

Catalog Number: RP1017-APC

About IL18

Interleukin-18 also known as IL18 is a protein which in humans is encoded by the IL18 gene. The protein encoded by this gene is a proinflammatory cytokine. IL-18 is a cytokine produced by macrophages and other cells that belongs to the IL-1 superfamily. IL-18 works by binding to the interleukin-18 receptor, and together with IL-12 it induces cell-mediated immunity following infection with microbial products like lipopolysaccharide (LPS). After stimulation with IL-18, natural killer (NK) cells and certain T cells release another important cytokine called interferon-gamma (IFN-gamma) or type II interferon that plays an important role in activating the macrophages or other cells. The combination of this cytokine and IL12 has been shown to inhibit IL4 dependent IgE and IgG1 production, and enhance IgG2a production in B cells. IL-18 binding protein (IL18BP) can specifically interact with this cytokine, and thus negatively regulate its biological activity. The human interleukin 18 gene IL18 maps to 11q22.2-q22.3, closely linked to the DRD2 gene locus and distinct from mapped IDDM loci.

Overview

Product Name	Anti-Interleukin-18 IL18 Antibody Picoband® APC Conjugated
Reactive Species	Mouse
Application	Recommended applications are based on the parent unconjugated antibody (ELISA, IHC, WB). Customers may select suitable applications according to their experimental needs.
Clonality	Polyclonal
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Rabbit
Uniprot ID	P70380

Technical Details

Immunogen	E. coli-derived mouse IL-18 recombinant protein (Position: N36-S192).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	APC Excitation Wavelength: 633-647 nm

	Emission Wavelength: 660 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

4 Publications Citing This Product

1. PubMed ID: 33681944, Yang L, Liu Y, Wang Y, Li J, Liu N. Azeliragon ameliorates Alzheimer's disease via the Janus tyrosine kinase and signal transducer and activator of transcription signaling pathway. Clinics (Sao Paulo). 2021 Mar 8;76:e2348. doi:10.6061/clinics/2021/e2348. PMID:33681944; PMCID:PMC7920406.
2. PubMed ID: 25049723, Si Lf, Zhang Sy, Gao Cs, Chen Si, Zhao J, Cheng Xc. Asian-Australas J Anim Sci. 2013 Oct;26(10):1399-405. Doi: 10.5713/Ajas.2013.13101. Effects Of Ifn-?? On Il-18 Expression In Pregnant Rats And Pregnancy Outcomes.
3. PubMed ID: 21920610, Wang Y, Zhang X, Zhang Y, Xu H, Fang G. J Reprod Immunol. 2011 Dec;92(1-2):45-53. Doi: 10.1016/J.Jri.2011.07.002. Epub 2011 Sep 13. Expression And Localization Of Il-18 In The Hypothalamic-Pituitary-Ovarian Axis Of Non-Pregnant, Pregnant, And Abor...

Visit bosterbio.com/anti-mouse-il-18-antibody-rp1017-boster.html to see all 4 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Interleukin-18 IL18 Antibody - APC

For Research Use Only. Not for use in diagnostic procedures.