

Anti-Interleukin-13 IL13 Antibody

Catalog Number: RP1047

About IL13

IL-13, Interleukin 13, is a protein that in humans is encoded by the IL-13 gene. IL-13 is a kind of cytokine secreted by many cell types, but especially T helper type 2 (Th2) cells, which is an important mediator of allergic inflammation and disease. The IL-13 gene is mapped to 5q23-q31. IL-13 induces its effects through a multi-subunit receptor that includes the alpha chain of the IL-4 receptor (IL-4Ralpha) and at least one of two known IL-13-specific binding chains. This gene acts more prominently as a molecular bridge linking allergic inflammatory cells to the non-immune cells in contact with them, thereby altering physiological function.

Overview

Product Name	Anti-Interleukin-13 IL13 Antibody
Reactive Species	Human
Description	Boster Bio Anti-Interleukin-13 <u>IL13 Antibody</u> catalog # RP1047. Tested in ELISA, WB applications. This antibody reacts with Human.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P35225

Technical Details

Immunogen	E. coli-derived human IL-13 recombinant protein (Position: L25-N146). Human IL-13 shares 57% and 62% amino acid (aa) sequence identity with mouse and rat IL-13, respectively.
Predicted Reactive Species	Bovine
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.



BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml, Human Direct ELISA, 0.1-0.5ug/ml, Human



Anti-Interleukin-13 IL13 Antibody (RP1047) Images



Boster Kit Box

2 Publications Citing This Product

1. PubMed ID: 24498391, Li J, Li L, Zuo H, Ke C, Yan B, Wen H, Zhang Y, Yang X. Plos One. 2014 Feb 3;9(2):E87887. Doi: 10.1371/Journal.Pone.0087887. Ecollection 2014. T-Helper Type-2 Contact Hypersensitivity Of Balb/C Mice Aggravated By Dibutyl Phthalate Via Long-Term De...

2. PubMed ID: 26771842, CD68 and interleukin 13, prospective immune markers for esophageal squamous cell carcinoma prognosis prediction

Visit <u>bosterbio.com/anti-il-13-antibody-rp1047-boster.html</u> to see all 2 publications.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Interleukin-13 IL13 Antibody